

Dec 22, 2017 Version 1

VASCULAR ENDOTHELIAL GROWTH FACTOR PLASMA LEVELS IN PATIENTS WITH CANCER OF THE LEFT COLON SUBMITTED TO OPEN OR LAPAROSCOPIC SURGERY V.1

DOI

dx.doi.org/10.17504/protocols.io.mdzc276

Marco Clementi¹, Mario Schietroma², Emanuele Lezoche³, Francesco Carlei², Alessio Agnifili⁴, Federico Sista², Sara Colozzi², Paola Mariani³, Francesco Masedu², Stefano Guadagni.²

¹Department of Life, Health and Environmental Sciences, University of L'Aquila, ITALY.;

²Department of Applied Clinical Sciences and Biotechnology, University of L'Aquila, ITALY;

³Department of General Surgery P. Stefanini, University of Rome "La Sapienza", ITALY;

⁴Department of Life, Health and Environmental Sciences, University of L'Aquila, ITALY



Marco Clementi

General Surgical Unit, San Salvatore Hospital, University of...

OPEN  ACCESS



DOI: dx.doi.org/10.17504/protocols.io.mdzc276

Protocol Citation: Marco Clementi, Mario Schietroma, Emanuele Lezoche, Francesco Carlei, Alessio Agnifili, Federico Sista, Sara Colozzi, Paola Mariani, Francesco Masedu, Stefano Guadagni. 2017. VASCULAR ENDOTHELIAL GROWTH FACTOR PLASMA LEVELS IN PATIENTS WITH CANCER OF THE LEFT COLON SUBMITTED TO OPEN OR LAPAROSCOPIC SURGERY. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.mdzc276>

Manuscript citation:

Unplished manuscript

License: This is an open access protocol distributed under the terms of the **Creative Commons Attribution License**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: December 22, 2017

Last Modified: March 22, 2018

Protocol Integer ID: 9369

Keywords: VEGF, open surgery, minimally invasive surgery, left colon cancer, angiogenesis.

Abstract

Background: VEGF is elevated in plasma of colon cancer patients and increases after surgery. This study's purpose was to determine plasma VEGF levels before, during, and after open and laparoscopic colorectal surgery performed for left colon cancer (CD).

Patients and methods: a multicenter, prospective, observational, non-randomized study was planned to evaluate samples obtained preoperatively, intraoperatively, and on postoperative days (1, 3, 7, 15, 30). VEGF-A₁₆₅ values were determined via ELISA. Statistical analysis was a repeated Anova with interactions.

Results: the recruitment finished in 4 years (71 patients). Baseline plasma VEGF-A₁₆₅ mean values (pVmv) resulted not significantly different in patients submitted to laparoscopy (125.4 pg/ml ± 47.7) and laparotomy (116.3 pg/ml ± 28.9). Intraoperatively, the highest difference was registered at the end of operations. In the 30-days postoperative period, the pVmv resulted significantly lower (G-G=0.45) in patients submitted to laparoscopy than in patients submitted to laparotomy (F= 144.6, p<0.001), at time went on (F= 97.6, p<0.001); the highest differences between groups were registered at day 7.

Conclusions: the trial supports the hypothesis that cancer and surgical trauma are related to production of VEGF-A₁₆₅ and demonstrates that open surgery, in comparison to laparoscopy, is a stronger promoter of VEGF release.

Attachments



[Main Manuscript prot...](#)

60KB



[Figures protocols.io...](#)

372KB

Guidelines

No guidelines

Safety warnings

- ⚠ Contradictory data regarding VEGF levels have been published; randomized prospective studies have not detected a significant difference on plasma and serum VEGF levels between laparoscopic and open surgery in colon cancer patients. However, it should be clearly noted (WARNING) that some reports had lack of homogeneity both in patient categories and surgical procedures.

1



